



DOI: **10.5958/2249-7137.2021.01019.3**

STUDY OF EPIDEMIOLOGICAL FEATURES OF VISCERAL LEISHMANIASIS IN ENDEMIC AREAS OF THE REPUBLIC OF UZBEKISTAN

Gaybullaev Fazliddin Khayriddinogli*

*Bukhara State Medical Institute named after Abu Ali Ibn Sino,
UZBEKISTAN

ABSTRACT

In the study of the clinical and epidemiological course of visceral leishmaniasis in the regions studied by the author, all 70 patients were children. The average age of the patients was 5.8 ± 0.8 . 58.6% (41 people) were girls and 41.4% (29) were boys. In terms of regions studied, the highest rate is in Navoi region - from 3.3 to 4.1 per 100,000 population, the incidence rate increased by 1.2 times, followed by Samarkand region - by 2.2 times. In Namangan region, there was a 2-fold decrease.

KEYWORDS: *Visceral leishmaniasis, sternal puncture.*

REFERENCES

1. Leishmaniasis: WHO bulletin. 2 March 2020 Link active on 20.09.20. Available at: <https://www.who.int/ru/news-room/fact-sheets/leishmaniasis>.
2. Obloqulov A.R., Narziev I.I., Kholov U.A., Niyazov G.E. (2018) Osobennostitecheniyakishechnogolyambliozauvzroslyx. New Day in Medicine 1 (21) 100-103;
3. Akhmedova M.D., Abidov Z.I., Gaybullaev F.X., Tashpulatova Sh.A., Anvarov J.A., Imamova I.A. Predictor of severe treatment of visceral leishmaniasis // NEWSLETTER OF THE MEDICAL ACADEMY OF MEDICINE OF ONTUSTIK KAZAKHSTAN - 2018. - P 36-41.]
4. Mirzoeva M.R. Character and frequency of complications in patients with chronic toxoplasmosis // New in medicine. - 2019. - C. 207-209.]

5. Gamal ShayfXaytemShayf Optimization of treatment of visceral leishmaniasis in children in Yemen // abstract on diss k.m.n.-Moscow.2006.-26s
6. Jirenkina E.N. Osobennostiochaga visceral leishmanioza in Papskomrayona, NamanganskoyoblastiUzbekistana // avtoreferatnadiss k.m.n.-Moskva.2013.-29s
7. MaxtiFaxar.
Sistematičeskijobzorimetaanalizdiagnostičeskoytestamochidlyadiagnostikivisistraln
ogoleyshmanioza.PROSPERO 2017<http://www.crd.york.fc.uk>
8. Chaturaka Rodrigo, Praveen Veratunga, SenakaRajapakse, Dipika Fernando.
Amphotericin For the treatment of visceral leishmaniasis: Systematic review and
metaanalysis Perspective guidelines for clinical trials, including the study of the
dosage of PROSPERO 2017<http://www.crd.york.fc.uk>.
9. Charles Abongomera, Saskia Van Xenten, Rezika Mohammed, Florian Fogt,
JozefenBuyze, ErmiasDiro, KoertRitmeyer, Tine Verdonk, Johann van Grevensven:
Systematic review of the factors of risk of death in patients with acute respiratory
infections. PROSPERO 2016<http://www.crd.york.fc.uk>