



DOI: **10.5958/2249-7137.2021.00833.8**

COMPARATIVE EVALUATION OF THERAPEUTIC EFFECT OF FERROCERONE IN ROSE SYRUP

Barno Inakova*

PhD, Docent,
Head of Department of Facultative Pediatric and Neonatology,
Andijan State Medical Institute, Andijan,
UZBEKISTAN

ABSTRACT

A new composition of ferroceron with improved organoleptic properties has been scientifically substantiated and proposed, which makes it possible to expand the scope of application of ferroceron in children with iron deficiency anemia, regardless of age. For the first time in young children with iron deficiency anemia, a comparative study of the therapeutic efficacy of ferroceron in rosehip syrup was carried out. At the same time, not only immediate but also long-term results of treatment were analyzed. The newly created composition of ferroceron in rosehip syrup, used as an iron-containing drug in the treatment of iron deficiency anemia in young children, turned out to be a highly effective agent without side properties. A comparative assessment of the antianemic effect of a combination of ferroceron and other iron-containing drugs, often used in children's practice, is given, which gave reason to recommend ferroceron with rosehip syrup as an antianemic drug that has an effect on the processes of hemoglobin synthesis and an erythrostimulating effect. The predominant both direct and long-term effect of treatment of children with ferroceron in rosehip syrup was revealed, which determined the created composition to be considered the drug of choice in the treatment of young children with iron deficiency anemia, an etiopathogenetic agent.

KEYWORDS: *Predominant, Syrup, Iron-Containing*

REFERENCES

1. Andrews N.C. Anemia of inflammation: the cytokine-pepsidin link //J.Clin.Invest.-2004.-V.113.-P.1251-1253
2. Antunes H., Goncales S., Teixeira-Pinto A. et all. Iron deficiency anemia in infants. Preliminary devolopment results at fivre years //Acta med.Port.-2005.-V.18.-#4.-P.261-266
3. Benefice E., Monroy S.L., Jimenez S., Lopez R. Nutritional status of Amerindian cyildren from the Beni River (Lewland Bolivia) as related to environ mental, maternal and dictory factors //Public Health.Nutr.-2006.-V.9.-#3.-P.327-335
4. Block J., Halliday J. Iron methobolism in health and diseas. W.B.Saunders company.-1994
5. Gurgoze M.K., Olcucu A., Aygun A.D. et all. Serum and nair levels of zinc, selenium, iron and copper wit iron deficiency anemia //Biol.Trace.Elem.Res.-2006.-V.111.-#1-3.-P.23-29
6. Herbert N., Giabol M.D., D.Suleymanova, Gregory W., Evons M.A. Anemia in young children of the Muynak district of Karakalpakstan, Uzbekistan. Prevelence, Tyne and correletes //Amer.J.of public Health.-1998.-V.88.-#5.-P.147-148
7. Hercberg S., Presiosi P., Galan P. Iron deficiency in Europe //Publ.Health.Nurt.-2001.-V.4.-#23.-P.537-545
8. Orellana J.D., Coimbra C.E., Lourenco A.E., Sentos R.V. Nutritional status and anemia in Surui Indian children, Brazilian Amazon //J.Pediatr. (Rio J.).-2006.-V.82.-#5.-P.383-388
9. Oliveira M.A., Osorio M.M., Rapaso M.C. Haemoglobin level and anemia in children in the state of Pernanbuco, Brazil: assosiation with socioeconomic and food consupcion factors //Cad.Saude Publica.-2006.-V.22.-#10.-P.2169-2178
10. Pierce M.B., Crowell R.E., Ferris A.M. Differing perspectives of inner-city parents and pediatric clinicians impact management of iron-deficiency anemia //J.Nutr.Educ.Behav.-2006.-V.38.-#3.-P.169-176